UNITED NATIONS

EP

UNEP(DEPI)/MED IG.22/Inf.11

UNITED NATIONS
ENVIRONMENT PROGRAMME
MEDITERRANEAN ACTION PLAN

21 December 2015
Original: English

19th Ordinary Meeting of the Contracting Parties to the
Convention for the Protection of the Marine Environment
and the Coastal Region of the Mediterranean and its Protocols

Athens, Greece, 9-12 February 2016

Agenda item 3: Thematic Decisions

Background document to the Regional Climate Change Adaptation Framework

For environmental and economic reasons, this document is printed in a limited number. Delegates are kindly requested to bring their copies to meetings and not to request additional copies.

UNEP/MAP
Athens, 2015

4.2.18 Spain 31
4.2.19 Syria 32
4.2.20 Tunisia 32
4.2.21 Turkey 32
4.3 International and Regional initiatives and frameworks 32
4.3.1 2WKHU 5HJLRQDO 6HDV ¶ $GDSWDWLRQ LQLWLDWLYHV 32
4.3.2 EU initiatives and policy frameworks 33
4.3.3 The Union for the Mediterranean (UfM) 34
4.3.4 Arab Framework Action Plan on Climate Change, 2010-2020 34
4.3.5 South East Europe 2020 Strategy 35
4.3.6 Global Adaptation Network 35
4.3.7 Mayors adapt 36
5 PRIORITY DIMENSIONS OF ADAPTATION POLICIES 37
5.1 Identifying and implementing low-regret measures 37
5.2 Integrating climate adaptation and Disaster Risk Management 38
5.2.1 Climate adaptation and Disaster Risk Reduction 38
5.2.2 The International DRR Framework 39
5.3 Integrating climate resilience into the ICZM process 39
5.3.1 Establishment stage 39
5.3.2 Analysis and Futures stage 40
5.3.3 Setting the Vision stage 40
5.3.4 Designing the Future stage 41
5.3.5 Realising the Vision stage 41
5.4 Ecosystems based Adaptation (EbA) 41
5.5 Promoting synergies between adaptation and mitigation 42
6 MOBILISING AND ACCESSING FUNDS FOR ADAPTATION 44
6.1 The international climate financing framework under the UNFCCC 44
6.1.1 Special Climate Change Fund (SCCF) 44
6.1.2 Adaptation Fund 44
6.1.3 Green Climate Fund (GCF) 45
6.2 Other international funding opportunities relevant to the Mediterranean region 45
6.3 Adaptation financing priorities at the national level 46
6.4 The role of the banking and insurance sectors 47
6.4.1 Enhance knowledge and raise awareness on climate risks 48
6.4.2 Reinforce climate risk prevention services 48
6.4.3 Development of insurance and/or investment products and services 48
6.4.4 Public sector role to create an enabling environment 49
7 BIBLIOGRAPHY 50

1 INTRODUCTION

1.1 Scope and structure of the report

The present report, submitted as an Information Document to the MAP Focal Points Meeting in Athens (13-16 October 2015 L V D E D F N J U R X Q G G R F X P H Q W D W L R Q W R W K H Regional Framework for Climate Change Adaptation Framework for the Mediterranean Marine and Coastal Areas, W Z D V S U H S D U H G in the context of the GEF-funded "Integration of climate variability and change into national strategies for the implementation of the ICZM Protocol in the Mediterranean" (ICZM) project which supported the finalization of the draft Framework.

human health and increasing inequality. Developing countries are particularly vulnerable because they are predominately poor, less resilient and lacking the capacity to manage disasters effectively.

The IPCC

- Actions to ensure that effective institutions, strategies, and choices for risk management will be identified, implemented, and sustained as an integrated part of achieving sustainable development.

The international community is increasingly rising up to the interdependent challenges of sustainable development, tackling climate change and risk management. 2015 is a critical year and a unique opportunity as it marks the conclusion of three international processes which will set the agenda through which the above issues are approached and addressed effectively and coherently in the years to come:

- In March 2015, at the Third World Conference on Disaster Risk Reduction in Sendai, Japan, UN Member States adopted the Sendai Framework for Disaster Risk Reduction (2015-30) which includes seven global targets and sets out four priority areas for further action. The new framework will guide countries in their efforts to achieve a substantial reduction of disaster losses in the future.
- By September 2015, governments will agree on a set of Sustainable Development Goals (SDGs) building on the outcome of the 2012 Rio+20 Conference and the Millennium Development Goals (MDGs). For the first time, these new goals will be designed for universal application.
- In December 2015, a comprehensive global agreement on climate change action is expected to be reached at the 21st Conference of the Parties to the United Nations Framework Convention on Climate Change (UNFCCC).

The following Table presents the three SDGs that are more relevant to adaptation to climate change in the Mediterranean (ie the ones on climate change itself, on the protection of seas and marine resources, and on the protection of terrestrial ecosystems), including the relevant targets, as they were agreed on 1 $XJXVW LQ WKH ILQDOL]HG WH[W IRU DGRSWLRQ ³7UDQVI 6XVWDLQDEOH 'HYHORSPHQW´

Sustainable Development Goal	Target	Target referring to specific means of implementation
Goal 13. Take urgent action to combat climate change and its impacts	13.1 Strengthen resilience and adaptive capacity to climate-related hazards and natural disasters in all countries 13.2 Integrate climate change measures into national policies, strategies and planning 13.3 Improve education, awareness-raising and human and institutional capacity on climate change mitigation, adaptation, impact reduction and early warning	13.a Implement the commitment undertaken by developed-

	14.7 By 2030, increase the economic benefits to Small Island developing States and least developed countries from the sustainable use of marine resources, including through sustainable management of fisheries, aquaculture and tourism	
Goal 15. Protect, restore and promote sustainable use of terrestrial ecosystems, sustainably manage forests, combat desertification, and halt and reverse land degradation and halt biodiversity loss	15.1 By 2020, ensure the conservation, restoration and sustainable use of terrestrial and inland freshwater ecosystems and their services, in particular forests, wetlands, mountains and drylands, in line with obligations under international agreements 15.2 By 2020, promote the implementation of sustainable management of all types of forests, halt deforestation, restore degraded forests and substantially increase afforestation and reforestation globally 15.3 By 2020, combat desertification, restore degraded land and soil, including land affected by desertification, drought and floods, and strive to achieve a land degradation-neutral world 15.4 By 2030, ensure the conservation of mountain ecosystems, including their biodiversity, in order to enhance their capacity to provide benefits that are essential for sustainable development 15.5 Take urgent and significant action to reduce the degradation of natural habitats, halt the loss of biodiversity and, by 2020, protect and prevent the extinction of threatened species 15.6 Ensure fair and equitable sharing of the benefits arising from the utilization of genetic resources and promote appropriate access to such resources 15.7 Take urgent action to end poaching and trafficking of protected species of flora and fauna and address both \demand and supply of illegal wildlife products	

maxima will increase not only because of an overall rise in mean temperature, but in some areas also because of increases in temperature variability and daily temperature range.

Precipitation is projected to decrease in all parts and all seasons (the most significant percent change occurs in summer) except for the northernmost parts in winter. The reduced summer rainfall is quite a homogeneous feature of the entire basin. During winter, however, the precipitation response appears to be different in the northern part of the Mediterranean area and over the Alps, where some increase of rainfall is found, whereas the rest of the domain characterized by a pronounced tendency to reduced precipitation.

In all of these simulations, the changes in mean seasonal precipitation during northern winter are characterized by a relative small decrease in the whole Mediterranean basin (in the range of -5%), with almost no change in the northern part of the basin and a more visible reduction (in the range of -5%-10%) in the southern part. During summer, all of the climate change projections show marked increase in the near-surface temperature and decrease in precipitation over the entire region.

Further, their results indicate high inter-model agreement on a significant decrease of soil moisture in all seasons, with the most significant decrease occurring in summer.

The CIRCE projections for the 21st century, therefore, suggest that remarkable changes in the climate of the Mediterranean region might occur already in the next few decades, showing a substantially steadily increase of the near-surface temperature that leads the Mediterranean lands to be about 2°C warmer in the 2021-2050 period with respect to the 1961-1990 mean. The change in the simulated near-

It is important to note that the steric contribution is only one of the components that might influence the sea level change in the Mediterranean Sea. There are other components that might determine the sea level trends in the basin, such as the melting of the continental ice sheets (Greenland and Antarctica) that, especially on the long term (centennial time scales), might become dominant.

It should also be noted that in the case of SLR in the Mediterranean, scientific uncertainty is particularly high, as making multi-decadal regional projections for relatively small isolated and semi-isolated basins such as the Mediterranean is more complex than for the global ocean (EEA, 2014). Nevertheless, the effect of SLR is considerable in most low-lying coasts of the Mediterranean basin where communities and infrastructure are typically located. In addition to the seawater expansion due to steric effect, coastal subsidence and global ocean level increase induced by continental glaciers melting (in Greenland and West Antartica) have to be considered as SLR components for the Mediterranean.

The following graphs are extracted from Annex I (Atlas of Global and Regional Climate Projections) of the Report of WGII in the IPCC AR5, which presents a series of figures showing for the



Figure 2: Time series for land and sea average temperature in the Mediterranean, under IPCC's Representative Concentration Pathway (RCP) scenarios. Source: IPCC, 2013: Annex I: Atlas of Global and Regional Climate Projections

The following table includes information on observed and projected regional changes in temperature and precipitation extremes, including dryness for the Mediterranean, extracted from Tables 3-2 and 3-3 of the SREX report. Projections are for the end of the 21st century versus the end of the 20th century and for the A2/A1B emissions scenarios.

	Trends in maximum temperatures	**Trends in**

3 OVERVIEW OF RISKS AND VULNERABILITIES

While determining tendencies and changes in the climatic system is quite delicate due to the multitude of factors that must be taken into account, the complexity of trying to identify the possible impacts of climate change is even greater, especially when considering uncertainties on regional and sub-regional trends. Indeed, these impacts are the result of confrontation between the major trends of climatic parameters and the specific conditions of the affected area, in other words the natural and manmade characteristics of the Mediterranean zone.

$V WKH ,3&& $5 Increasing magnitudes of warming increase the likelihood of severe, SHUYDVLYH DQG LUUHYHUVLEOH LPSDFWV >«@ *OREDO FOLPD
mean temperature increase of 4°C or more above preindustrial levels in all reasons fo

The changes of climate patterns will result in various physical impacts on water resources, since water is involved in all components of the climate system, and affects different aspects of human well-being from agricultural productivity and irrigation supply to flood and drought control, municipal and

3.1.3 Coastal systems and low-lying areas

Coastal zones, arguably the most appealing assets of the Mediterranean, are already exposed to significant pressures from land-based and marine pollution, urban development, fishing, aquaculture, tourism, damming, extraction of materials, and marine biological invasions. Climate change, and especially the major driver of sea level rise, is expected to significantly increase these pressures. In particular, many coastal systems will experience increased inundations and storm flooding, accelerated coastal erosion, seawater contamination of fresh groundwater, displacement of coastal lowlands and wetlands, encroachment of tidal waters into estuaries and river systems, possible loss of nesting beaches. More frequent and severe weather and climatic events will further enhance these phenomena, while in the longer term, changes in wind and wave patterns could interfere with sediment transport leading to greater erosion or accretion.

Coastal erosion will lead over time to the inland migration of the beaches of the Mediterranean with soft sedimentary coasts being more vulnerable than harder, rocky coastlines. River deltas, due to their particular topography, are particularly vulnerable to the impacts of erosion and inundation. Damming of rivers upstream no longer allows the normal circulation of sediment, which cannot reach the delta to consolidate it.

At the local scale, possible impacts from sea level rise are also determined by other non-climatic factors such as the subsidence of coastal land, subsurface resource extraction, and tectonic movements.

The JRC PESETA II Project ³&OLPDWH ,PSDFWV LQ (XURSHºC scenario the HVWLPDWHV W average annual costs from sea floods damage in Southern Europe will increase from 163 to 903 million in the 2080s.

3.1.4 Ocean systems

The Mediterranean Sea is among the richest in biodiversity of global importance, rich with endemism and autochthonous species. At the same time, it has unique marine features that make this region particularly vulnerable to climate change. The overall extent of water exchange is restricted due to the narrow connections with the Atlantic Ocean, the Red Sea and the Black Sea. In addition, due to the relatively small size of the basin, seawater in the Mediterranean can more easily heat up and evaporate, combined with hot, dry summers and low inflow from rivers.

Increases in sea temperatures will alter distribution of species and foster the spread of warm water species into the Mediterranean, thus promoting the displacement of ecotypes and shifts in ecosystem functioning and ultimately lead to loss of species. The IPCC AR5 identified the Mediterranean Sea as one of the semi-enclosed seas with projected high rates of local extinction because land boundaries will make it difficult for species to move laterally to escape waters that may be too warm. Additionally, periods of extreme seawater temperature during heatwaves will contribute to mortality events that affect many invertebrate species as well as Posidonia meadows.

Another emerging climate-related threat to Mediterranean marine ecosystems, is ocean acidification, the phenomenon of shifting the chemical balance of seawater to a more acidic state (lower pH) due to increased CO2 concentrations in the sea as a result of increased CO2 concentrations in the atmosphere. Acidification is currently occurring at a geologically unprecedented rate, subjecting marine organisms to additional environmental stresses. According to

Mediterranean seawater has increased by 10% since 1995 and if current CO2 emission rates continue, it will increase another 30% by 2050 and 150% by 2100. Several planktonic organisms are affected by acidification with possible negative impacts on fish populations. Moreover, acidification also threatens iconic and invaluable Mediterranean ecosystem-building species (such as sea grass meadows, Coralligene reefs and Vermetid snail reefs) which create rich key habitats and homes to thousands of species, and also protect shores from erosion as well as offer a source of food and natural products to society.

More specifically, the following Mediterranean ecosystem types will be affected by climate change: [source: MedPAN 2012]

Coralligenous formations: Sea warming and exposure to high temperatures may affect coralligenous formations both directly, by causing harmful metabolic changes and physiological stress, and indirectly by reducing their resistance to pathogens. Heatwave events involving a 3 – 4 °C increase in seawater temperature can induce mass-mortality events in some gorgonian or sponge communities. Coralligenous species with calcareous skeletons can be affected by increased seawater acidification, which both reduces skeletal growth and tends to dissolve their skeletons. As most of these species are characterized by high longevity and slow growth, a dramatic increase in mortality could have severe consequences in terms of population viability, and consequently could affect the structure and function of coralligenous formations as habitat-forming assemblages in the marine ecosystem.

Coastal fish populations: Climate change is expected to force fish species distributions toward the northern areas of the Mediterranean, especially for coastal fishes. Species typical of the southern parts of the Mediterranean, are now spreading northward into the coldest parts of the Mediterranean as a result of rapid warming. Other thermophilic fishes have increased in abundance with the rise in water temperature. Conversely, cold-water species are contracting their distribution ranges as a consequence of warming and, in the worst-case scenario they could eventually face extinction. The warming of the

3.1.5

3.2 Human Settlements, Industry, and Infrastructure

3.2.1 Urban areas

More than a third of the population of Mediterranean countries live in coastal zones. This is more evident in Northern African and Middle Eastern countries where coastal cities have traditionally been particularly important due to the aridity of inland regions.

As coastal populations and assets in coastal areas continue to grow, exposure to climate change-related hazards ±and especially those associated with sea-level rise- is also increasing. What is especially worrying is the rapid growth of highly vulnerable urban communities living in informal settlements, many of which are on land at high risk from extreme weather and often lack essential infrastructure and

increase in the influx of tourists to the coast in the months when the air and water temperature will not be too hot, thus shifting from the hot summer months to the spring and autumn months.

However, there is considerable uncertainty about how tourists will respond to the effects of climate change and therefore overall vulnerability of coastal tourism is hard to assess.

Transportation

Port infrastructure, but also coastal roads, railways, and airports, are expected to be at risk mainly due to temporary and permanent flooding arising from sea-level rise, high winds and storm surges. Negative

in mortality is high: total deaths from natural causes increased by 14%, deaths from cardiovascular problems by 22% and respiratory problems by 32% during heatwaves events. According to recent

-

The application of the CRI-MED methodology to the eleven Mediterranean countries selected for the project ClimVar & ICZM led to a ranking of the relative risk of each coastal region in relation to potential coastal hazards (coastal erosion, coastal flooding and saltwater intrusion).

The distribution of coastal segments to the 5 risk classes is shown in the following table and map:

Risk class	**CRI Score**	**Number of cells**	**% of the total**
Extremely High	0,55 -		

change-related issues in planning for sustainable adaptation and ICZM strategies on their national coastal territory.

1996	2001	2005	2007	2010	2015
Observing					

Egypt is also a member of the Nile Basin Initiative, a partnership among states along the Nile Rivers established to encourage sustainable socioeconomic development through the equitable division of the 1LOH %DVLQ¶V ZDWHU UHVRXUFHV. The initiative has begun to address climate change within this regional IUDPHZRUN DQG LQ ODXQFKHG WKH SURMHFW ³$GDSWLQJ WR 1LOH 5LYHU %DVLQ´ ZLWK DVVLVWDQFH IURP 81(3 DQG WKH 6 (SIDA).

4.2.7 France

In France, a National Adaptation Strategy was adopted in 2006. Regional impact, vulnerability and adaptation studies were carried out to implement subnational adaptation planning policies (Regional Climate, Air and Energy Schemes process).

A national adaptation plan (NAP) was adopted in 2011, with a five-year implementation period. A review of actions on NAP implementation took place in June 2013. The NAP has prioritised 240 concrete measures covering the 20 thematic areas of the plan: Cross-cutting actions, Health including "Plan Canicule", Water, Biodiversity, Natural hazards, Agriculture, Forests, Fisheries and aquaculture, Energy and industry, Transport infrastructures, Urban planning and the built environment, Tourism, Information, Education and training, Research, Funding and insurance, Coastlines, Mountains, European and international actions, Governance.

4.2.8 Greece

Adaptation policy progress in Greece is still in the agenda-setting stage.

The Ministry of Reconstruction of Production, Environment and Energy, the Athens Academy and the Bank of Greece signed a memorandum of cooperation in the implementation of a "National Strategy for Adapting to Climate Change". The aim of their cooperation is to create a mechanism to monitor implementation of the national strategy and incorporate adaptation policies in all areas. The National Strategy for Adapting to Climate Change is being developed and it will be completed by the end of 2015.

In March 2009, on the initiative of the Bank of Greece, a Committee of scientists was set up in order to produce a study of the environmental, economic and social impacts of climate change in Greece. The Committee finalised the first phase of the project and in June 2011 published a comprehensive study of the impacts of climate change in Greece, addressing in particular the cost of climate change for the Greek economy, the cost of implementing adaptation measures, and the cost of moving to a low-emissions economy in the context of EU climate change mitigation policies. In the current second phase of the project, the Committee is working on a new study to contribute to the formulation of a national adaptation strategy on climate change.

4.2.9 Israel

In 2011, the Ministry of Environmental Protection set up the Israeli Climate Change Information Center, which aims to develop the scientific knowledge base and policy documents that will feed into the national adaptation plan. The Centre has since submitted three reports ±the first, in 2012, reviewed existing knowledge on the issue, and identified and prioritised knowledge gaps. The second, also in 2012, provided policy recommendations and an international marketing programme for ICCIC deliverables, while the third, in 2013, reviewed adaptation to climate change in local authorities.

4.2.15 Morocco

Morocco has developed a National Plan against Global Warming (PNRC) that was presented at the COP 15 held in Copenhagen in 2009. The Plan provides for reducing greenhouse gas emissions through the development and diversification of clean energy sources and the implementation of adaptation measures that rely mainly on the water strategy and Green Morocco Plan for Agriculture, also launched in 2009.

$ ZLGH UDQJH RI DGDSWDWLRQ WRROV KDYH EHHQ LQFRUSRU
such as in the Water Sector, Agriculture, Forestry, biodiversity and combating desertification, Housing,

Environment on 2012. The Framework aims at enhancing th H $U D E &R X Q W U L H V ¶ F D S D F appropriate measures for addressing climate change issues while achieving sustainable development targets and MDGs in the Arab Region. It includes mitigation and adaptation objectives. UNEP ROWA and UN ESCWA provided technical assistance for its elaboration.

Technology Transfer and Climate Change Action (REGATTA) in Latin America and the Caribbean, the

5.2.2 The International DRR Framework

On the 3rd United Nations World Conference on Disaster Risk Reduction (WCDRR) in Sendai, Japan (14-18 March 2015), more than 5,000 government officials, ministers and leaders from over 160 countries agreed a new global framework to guide decision makers towards a more disaster-resilient future, known as the Sendai Framework.

The agreed text places a strong emphasis on tackling the underlying drivers of risk, including: poverty, climate change, ecosystem decline, bad urban planning, land use and risk governance. It includes seven global targets (most of which are not quantifiable) and sets out four priority areas for further action, presented below:

Targets

- Substantially reduce global disaster mortality
- Substantially reduce the number of affected people globally
- Reduce direct disaster economic loss in relation to global GDP
- Substantially reduce disaster damage to critical infrastructure and disruption of basic services
- Substantially increase the number of countries with national and local disaster risk reduction strategies
- Substantially enhance international cooperation to developing countries
-

These guidelines propose three types of indicators:
1.

However, it is crucial to avoid negative trade-offs between adaptation and mitigation, or between economic goals and environmental goals.

In some cases, for example, adaptation measures may increase greenhouse gas emissions. Such examples include the increase in the consumption for fossil fuels for air conditioning in response to higher temperatures or desalination and increased water pumping. Similarly, some mitigation measures may impede adaptation, such as the construction of dams for hydropower which contribute to the reduction of sediment input into the sea and therefore aggravate coastal erosion, or the production of biofules which may increase competition for limited land and water resources.

The Adaptation Fund was established under the Kyoto Protocol and has been financed primarily through a 2% levy on proceeds from the Clean Development Mechanism (CDM), supplemented by voluntary contributions. It supports concrete adaptation projects and programs in developing countries that are particularly vulnerable to the adverse effects of climate change. The fund is operated by the Adaptation Fund Board, which has dedicated $265 million to increase climate resilience in 45 countries. However, with demand and prices for CDM credits currently at very low levels, Annex I parties and international RUJDQL]DWLRQV KDYH EHHQ HQFRXUDJHG WR VFDOH XS IXQG fundraising goals of $80 million per year in 2014 and 2015.

6.1.3 Green Climate Fund (GCF)

The Green Climate Fund, established under the COP 16 Cancun Agreements in 2010, is expected to become the main instrument to finance adaptation measures in much of the developing world. With the initial resource mobilization process of the Fund being completed at the end of 2014 with pledges of 10.2 million dollars, the GCF is scheduled to start programing resources in the second half of 2015. The GCF Board has decided that 50% of its portfolio should be allocated to adaptation and, in turn, that 50% should go to particularly vulnerable developing countries including Least Developed Countries (LDCs),

- the Western Balkan Investment Framework and
- the Islamic Development Bank.

In addition to these major multilateral funds, many developed countries also finance adaptation in self-selected developing countries through their respective bilateral development assistance agencies. Key bilateral donors in the agenda include France, Finland, Germany, Italy, Netherlands, Spain, Sweden as

the focus is on flooding which is established as a natural phenomenon exacerbated by climate change. The project consists of three phases, the first of which ±a report on how to build disaster-resilient communities ±was launched in June 2014. The second phase comprises the building of global and country natural disaster risk maps to identify partic

Responsible and effective property and agriculture insurance companies should fully integrate natural disaster exposure zones in their practices.
At the regional level, the options for establishing a Mediterranean Catastrophe Risk Insurance Facility could be explored, taking into account the experience of the relevant Facility for the Caribbean. Regional risk management institutions and platforms could be established to foster collaborative action to coordinate the management of weather-related risks and build risk prevention and management capabilities.

6.4.4 Public sector role to create

ESCWA, 2011. Assessing the

List of Abbreviations

CAF

SREX Special Report on Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation
SST Sea Surface Temperature
UfM Union for the Mediterranean
UN ESCWA United Nations Economic and Social Commission for Western Asia
UNCLOS United Nations Convention on the Law of the Sea
UNEP United Nations Environment Programme
UNEP ROWA UNEP Regional Office for West Asia
UNFCCC United Nations Framework Convention on Climate Change
WCRP World Climate Research Programme

Glossary of Terms

Adapted from: IPCC, 2014. Climate Change 2014: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.
ICZM definition from the Protocol on Integrated Coastal Zone Management in the Mediterranean.

Adaptation:	The process of adjustment to actual or expected climate and its effects. In human systems, adaptation seeks to moderate or avoid harm or exploit beneficial opportunities. In some natural systems, human intervention may facilitate adjustment to expected climate and its effects.
Climate change	Climate change refers to a change in the state of the climate that can be identified (e.g., by using statistical tests) by changes in the mean and/or the variability of its properties, and that persists for an extended period, typically decades or longer. Climate change may be due to natural internal processes or external forcings such as modulations of the solar cycles, volcanic eruptions, and persistent anthropogenic changes in the composition of the atmosphere or in land use.
Climate-resilient pathway for development	A continuing process for managing changes in the climate and other driving

	interactions, the maritime orientation of certain activities and uses and their impact on both the marine and land parts.
Impacts:	