Economic & Social Affairs

United Nations Expert Group Meeting on Current Issues in the Estimation of Adult Mortality

New York, 26-27 October 2006

Report of the Meeting

United Nations

DESA

The Department of Economic and Social Affairs of the United Nations Secretariat is a vital interface between global policies in the economic, social and environmental spheres and national action. The Department wo

PREFACE

The Population Division of the Department of Economic and Social Affairs of the United Nations Secretariat is in charge of estimating levels and trends of mortality for all the countries of the world. The work of the Population Division in this area has encompassed both the preparation of estimates of mortality indicators and the development of methods to estimate mortality, particularly when the data available are incomplete or deficient. The most recent contribution of the Population Division in the area of methodology is the report entitled *Methods for Estimating Adult Mortality*,[1] which presents the major approaches to the estimation of adult mortality using indirect or incomplete information.

The Population Division has also contributed to the development of mortality models that may be used in conjunction with available evidence to derive full life tables for populations lacking complete and reliable information on deaths by age. The United Nations model life tables for developing countries[2] provide a set of models that were derived in the early 1980s on the basis of the mortality experience of developing countries with reliable information.

With the establishment of the Mortality Section in 2004, the Population Division is devoting more attention to the improvement of both the methods used to estimate mortality and the development of flexible models allowing the derivation of life tables from partial information. This work benefits from a dialogue with experts engaged in similar work both within and outside the United Nations system. The Expert Group Meeting on Adult Mortality convened by the Population Division provided an opportunity to discuss ongoing activities in other institutions and to obtain feedback on the work carried out by the Population Division. As mortality levels continue to decrease in most countries of the world, deaths become increasingly concentrated at older ages and it becomes more pressing to estimate mortality in adult ages with accuracy. In addition, as the AIDS epidemic continues to expand, accounting for its effect on adult mortality is essential. These are some of the issues discussed by the Expert Group Meeting, as reflected in this report.

This report as well as other population information can be accessed via the Internet on the official website of the Population Division, www.unpopulation.org. For further information concerning this publication, please contact the office of Ms. Hania Zlotnik, Director, Population Division, Department of Economic and Social Affairs, United Nations, New York, NY 10017,

Explanatory Notes

The following acronyms are used in the report:

DDM	Death Distribution Methods
DHS	Demographic and Health Surveys
DSP	Disease Surveillance Points
GGB	General Growth Balance
HIV/AIDS	Human Immunodeficiency Virus/Acquired Immunodeficiency Syndrome
INED	Institut national d'études démographiques
INDEPTH	International Network for the continuous Demographic Evaluation of Populations and their Health in developing countries
MDG	Millennium Development Goals
MICS	Multiple Indicator Cluster Surveys
PAHO	Pan American Health Organization
SEG	Synthetic Extinct Generations
SRS	Sample Registration System
UNAIDS	Joint United Nations Programme on HIV/AIDS
UNFPA	United Nations Population Fund
UNICEF	United Nations Children's Fund
WHO	World Health Organization

UNITED NATIONS EXPERT GROUP MEETING ON CURRENT ISSUES IN THE ESTIMATION OF ADULT MORTALITY

A. OPENING OF THE MEETING

Ms. Hania Zlotnik, Director of the Population Division, opened the meeting by welcoming participants and noting that the topic they would discuss had particular relevance in the context of recent trends and developments in the areas of health and mortality. She pointed out that there was an increasing need for reliable estimates of adult mortality, particularly for developing countries. These estimates would allow to establish whether, as was expected in the 1970s or early 1980s, the populations of developing countries would undergo the epidemiological transition as those of developed countries had. She recalled that, according to the epidemiological transition, mortality decline would be accompanied by a reduction of mortality caused by infectious disease and a predominance of deaths caused by chronic or degenerative diseases. The advent of HIV/AIDS and its rapid spread in sub-Saharan Africa already meant that the countries most affected by the epidemic were not experiencing a reduction of mortality due to infectious diseases but adequate estimates of adult mortality for those countries were largely lacking. Furthermore, recent research had shown that some diseases once considered degenerative, such as liver, stomach or cervical cancer, were actually caused by infectious agents. It was an open question whether the spread of those diseases would be controlled rapidly enough through relatively low-cost interventions such as vaccines. Recent evidence suggested that the prevalence of diabetes was increasing in many developing countries. Whether their health systems would be capable of providing the treatments necessary to extend the life of those affected was another important question. Better estimates of adult mortality that allowed tracking changes over time would facilitate planning and decision-making regarding the type of interventions that were necessary or most effective in different contexts.

C. R

sus of the population living in the site. Some of the surveillance sites had been operational for a long time, such as Matlab in Bangladesh, which had been in operation for 40 years. The INDEPTH network was established in 1998.

It was possible to pool data from multiple sites to make estimates of mortality. In 2002, the Network had published a monograph on mortality and was in the process of updating that work. It had also produced a monograph that presented new model life tables derived from the experience in the surveillance sites and a journal article on causes of death in Africa. Mr. Bawah recognized that the estimates based on surveillance data were not nationally representative and said that INDEPTH was starting to work with the national statistical offices of some countries to try and explore ways of obtaining better information at the national level. Countries having several surveillance sites might obtain reasonably good national estimates from their pooled data, although whether the sites in existence were adequately representative of the whole population would remain an issue.

INDEPTH was also collaborating with the Health Metrics Network and with WHO on a new programme to study adult mortality. Based on the voluntary collaboration of sites run by different institutions whose objectives differed, INDEPTH was facing the problem of ensuring that all sites made their data available to the network.

University of California at Berkeley

Ms. Dana Glei, Research Demographer of the University of California at Berkeley, reported that the Human Mortality Database, a joint project of the University of Berkeley and the Max Planck Institute for Demographic Research, contained time series data on mortality for 28 national populations with high-quality data. The project was funded by the U.S. National Institute on Aging until 2009. The guiding principles in developing the database were to ensure the comparability of the data across countries and over time; to maintain flexibility by allowing the data to be retrieved in a variety of formats; to ensure accessibility to both the estimates derived from the data and to the basic data themselves; and to ensure that the results obtained could be reproduced by any independent researcher. In obtaining, assessing and cleaning the basic data, the project relied on national expertise. While the database did not include any developing country, the project had provided some support to another University of California project focusing on Adult Mortality in Developing Countries and was supporting a researcher at Rockefeller University who was focusing on the analysis of data for developing countries.

Harvard University

Mr. Ken Hill, Professor at the Harvard Center for Population and Development Studies, reported that the Center was testing data collection instruments, including a number of new questions on mortality and cause of death, in four field sites. One objective was to develop a standard method to conduct verbal autopsies. The Center was engaged in carrying out several studies of programme evaluation of health operations in a number of countries and a new project had been initiated in Mexico. There was also interest to continue work in testing the performance of indirect methods for the estimation of adult mortality. A proposal to continue the project on Adult Mortality in Developing Countries was under review. Lastly, the Center would collaborate in revising the Global Burden of Disease estimates which, aside from estimates for the major world regions would also include country-specific estimates for about 15 countries.

D. CURRENT ESTIMATES OF ADULT AND OVERALL MORTALITY

Ms. Mie Inoue, Statistician at WHO, presented WHO current practice regarding data and methods to construct life tables. Empirical data were compiled from vital registration, sample registration systems, demographic and health surveys, and population censuses. WHO requested that its Member States annually submit data on deaths classified by age, sex and cause. The completeness of death registration was assessed by using demographic techniques such as the

Ms. Inoue suggested that, in order to harmonize WHO estimates of life expectancy at birth with those prepared by the United Nations Population Division, one would need to focus first on using similar estimates of the completeness of death registration and the same version of UNAIDS estimates. She welcomed the idea of convening meetings such as this one to discuss adult mortality estimation and facilitate data sharing and the assessment of completeness of vital registration.

Ms. Cheryl Sawyer, Population Affairs Officer at the Population Division, presented the data and methods used by the United Nations Population Division to estimate mortality in preparing the revisions of *World Population Prospects*, the Division's biennial assessment of past population trends and future prospects. The Population Division's projection software, Abacus, had been in use since the 1970s but had been modified in significant ways, particularly with respect to the procedures used to project mortality. Abacus projected populations classified by five-year age groups with an open-ended interval of 100+ over five-year periods. Calculations started at a base year, that could range from 1950 to 1995 in the *2004 Revision*. Empirical information on the components of population change was used to project the population forward from the base year until 2005 and, from there on, models were used to project the population in the future.

Regarding mortality, the empirical information examined by the Population Division in developing its estimates included some combination of the following: (*a*) life expectancy at birth; (*b*) under-five and infant mortality rates (derived in collaboration with the Inter-Agency Group for Mortality Estimation); (*c*) age-specific mortality rates derived from vital registration data adjusted as necessary; (*d*) life tables calculated by other institutions or researchers; (*e*) indirect estimates of various kinds, and (*f*) estimates of HIV prevalence obtained from UNAIDS. The mortality inputs for the projection program could be provided in the form of a set of survival ratios by sex (s_x) for each period, that is, the proportion of individuals in a five-year age group ($_5L_x$) who would be alive five years later ($_5L_{x+5}$) according to the age-specific mortality rates of that period; or the level of life expectancy by sex and the selection of a family of model life tables from which the necessary survivorship ratios would be derived. There was also the option of providing the survival ratios by sex for the base or initial period, and levels of life expectancy by sex for subsequent periods (the so called "modified method"). This option was used for most countries with adequate empirical data on mortality by age and sex because it permitted the automatic generation of appropriate survivorship ratios consistent with the expectation of life provided as input for each period and with the age pattern of mortality in the initial survivorship ratios modified, over time, according to the rates of change of the survivorship ratios in an underlying family of model life tables.

For countries lacking information on age-specific mortality, input took the form of estimates of life expectancy by sex and a model life-table family selected to provide the age pattern of mortality. When only the level of child mortality was known, the estimated life expectancy was derived using a family of model life tables.

Ms. Sawyer noted that the mortality estimates used in estimating population trends for past periods were an integral part of a comprehensive and consistent set of population estimates and the need to ensure consistency with enumerated populations was sometimes the cause of discrepancies between the estimates produced by the Population Division and those obtained while focusing only on mortality data. If a new system to generate life tables and survivorship ratios from existing evidence were to be developed, it would have to be added to the options available to project mortality in Abacus and would likely take some time to become operational.

Mr. Kirill Andreev, Population Affairs Officer at the Population Division, presented comparisons of current mortality estimates made by WHO and the Population Division. For half the countries, the estimated life expectancy at birth in the two sources differed by a year or less, but for a quarter it differed by at least two years. Regarding adult mortality or, more specifically, the probability of dying between exact

but one, the information available was deficient or incomplete. However, most countries had sufficient information for the estimation of at least mortality in childhood and a high proportion also had information allowing the estimation of some indicator of adult mortality. Hence, the aim was to provide a means

There was interest in knowing to what extent the parameter *k* could be used to reflect the special situation in a country, even if a reliable estimate of adult mortality were not available. For instance, could *k* be used to reflect the impact of HIV/AIDS? There was also interest in exploring ways of incorporating some measure of uncertainty when using the model to generate mortality schedules.

Regarding the assessment made by Mr. Canudas-Romo, it was observed that it was not very revealing to analyse the fit to e_0 and ${}_{45}q_{15}$ separately, since *e*

lated data affected by different types of errors were used in the application of the two methods. Four different types of errors were simulated: (*a*) a completeness of enumeration that varied with age; (*b*) a completeness of death registration that varied with age; (*c*) age misreporting, and (*d*) non-zero net migration.

According to the tests performed, age misreporting did not have a serious effect on the estimates yielded by either of the two methods. Non-zero net migration, in contrast, had a noticeable effect on the results produced by both methods and the effect was more marked the larger the magnitude of net migration, with the results of the SEG method being more affected than those of the GGB method. Regarding a completeness of enumeration that varied with age, it had a larger effect on the estimates produced by the GGB method than on those yielded by the SEG method, with the size of the effect related to the variance of completeness of enumeration by age. A completeness of death registration that varied with age had effects that varied from modest to substantial. In addition, the results of the SEG method were sensitive to changes in the completeness of enumeration from one census to the next. In view of these results, it was suggested that using both methods in combination could yield better estimates than using either of them separately. The best strategy was to use the GGB method to derive an adjustment for changes in the completeness of enumeration from one census to the next, adjust the second census on that basis and then use the adjusted results in the application of the SEG method to obtain an estimate of the completeness of death registration.

Ms. Lioudmila Andreeva, Research Specialist at Rockefeller University, reported on work she was carrying out to update the data for developed countries contained in the Human Mortality Database and to check the consistency of data on population and deaths by age for developing countries as contained in various secondary sources, including the United Nations Demographic Yearbook database, the WHO mortality database, and the data compiled for the Adult Mortality in Developing Countries project of the University of California at Berkeley. In checking the information available in these sources, she had found many inconsistencies among data that were supposed to originate from the same source but whose versions in the different secondary sources considered did not always coincide. Consistency among the different secondary sources improved the more recent the data considered. Because the secondary sources used did not have adequate information on the original sources from which the data had been obtained, it was not always possible to establish the reason for the inconsistencies detected. Ms. Andreeva noted that, in planning a database containing the basic data for mortality estimation, attention had to be given to the type of meta-information that it would be useful to include regarding the original data sources used and the types of adjustments that the data compiled in the database had been subject to. She was sharing the results of her work with the Population Division.

Mr. Wilmoth discussed a strategy to develop a protocol for compiling, organizing and analysing the data required for the application of death distribution methods. The protocol would include a description of the various formats in which the original data appeared; means of transforming the original data into standardized formats; details of how the estimation methods would be applied; and the types of checks, including diagnostic plots, to which the results of the estimation methods would be subject. Examples of what he considered format issues included: how to treat vital registration series with missing data for a year or a number of years; how to treat residual categories, such as those labelled “unknown” or “not stated”; how to handle non-standard age groups; or how to deal with differences in the open-ended age group. Developing a comprehensive protocol on how to deal with any of these contingencies would ensure consistent preparation of the data for use with the death distribution methods and would facilitate their consistent application across populations and over time, thus permitting a fair comparison of the results obtained. It would also facilitate developing appropriate software to automate the updating of mortality indicators. Mr. Wilmoth considered that the preparation of such a protocol was a necessary and useful basis for inter-agency collaboration in sharing information and data for the estimation of adult mortality.

The discussion considered first whether the death distribution methods could be adapted to include explicit allowance for international migration. If reliable independent estimates of net migration by age were available, adjustment was possible. Unfortunately, very often the best estimates of net migration

Mr. Andreev said that all incoming data would be archived in their original format and listed in a reference database. The data would be evaluated and reformatted, as necessary, to be stored in a database, tentatively being called "Database of Demographic Quantities" which would be a repository for all demographic data useful for mortality estimation. The database would also be used for the dissemination of mortality estimates and it would be developed in a relational model compatible with existing databases in the Population Division.

Mr. Andreev presented an overview of the relational database model proposed. Demographic quantities, including deaths by age and sex, population by age and sex, data on children ever born and surviving, and official life tables, would be input and each of the data items would be qualified in terms of a reference date (calendar time), age, sex, Lexis shape and other relevant variables. To implement the different methods of mortality estimation, appropriate software would be developed as a library in Matlab. To ensure quality control, checking programs would be developed.

Ms. Sawyer described the inventory of all available primary data for mortality estimation that was being carried out in preparation for the development of the database. A database on data availability had been set up and had been tested. It allowed the production of reports on the availability of data by country or data type. This database would eventually be part of the Database on Demographic Quantities and would guide the development of the latter. To populate the inventory database, the census questionnaires from the 1990 and 2000 rounds of population censuses had been reviewed for all content relevant to the estimation of mortality. The Population Division's data files on child mortality estimation had been consolidated and updated to produce a comprehensive list of population surveys and censuses that contained information allowing the estimation of child mortality. Work on compiling the sources and the type of data relevant for the estimation of adult mortality was starting. The collaboration of partners in identifying all possible data sources would be appreciated. Mr. Andreev presented an Excel worksheet that collaborators could use to submit the required information.

Mr. Wilmoth proposed a path for collaboration in developing the mortality database. The main steps in carrying out this task included: (*a*) designing the database; (*b*) constructing the database; (*c*) populating the database, and (*d*) maintaining the database. Along the way, it was important to ensure access to all collaborators, to maintain quality control, and to develop the necessary analytical features or capabilities. It was part of the work programme of the Population Division to improve mortality estimation and the database was a key instrument to do so. Hence, the Population Division was in the process of developing the database but, because other institutions already had considerable experience with similar databases, it was seeking their collaboration. Such collaboration would reduce duplication of work and potentiate available resources. To facilitate collaboration, the plan was to develop a web-based interface to allow interested partners to contribute by populating the database and to have access to its contents. During the development phase, access would be restricted to collaborating institutions but it was expected that, eventually, most of the contents of the database would be publicly available. Regarding the development of analytical tools for use in connection with the database, the first set would likely include tools to assess data quality and make data comparisons. During a second stage, tools allowing the application of estimation methods or the evaluation of their results would be developed. He then outlined some of the future steps: (*a*) complete the database design and develop the database; (*b*) define a collaborative strategy to populate the database; (*c*) develop the necessary queries and analytical features for the web-based use of the database, and (*d*) open the database for use by the collaborating institutions.

During the discussion, participants offered suggestions about the type of data that should be gathered. With respect to data relevant for the estimation of child mortality, for instance, it was suggested that, in addition to the number of children ever born and surviving by age of mother, the full tabulations of women classified by number of children ever born and women by number of children surviving (or dead) should also be included because they allowed additional checks of data quality. It was noted that to

gather more detailed data than those generally published, it would be necessary to obtain special tabulations from certain sources. One example involved data by sex on children ever born and surviving, which were not normally published in the reports of the Demographic and Health Surveys but could be obtained from the survey data themselves.

Participants suggested that, to the extent possible, it would be useful to automate the input of data into the database, especially when the primary data were already available in electronic form. Once more, the need to record the necessary meta-information on all data was underscored. Allowance could be made, for instance, to record the notes accompanying published tables. Regarding the treatment of missing data, it was suggested that allowance be made for the recording of any information available on whether and what type of imputation had been made. It might also be useful to code the results of the qualitative analysis of each data set to make them transparent to users. In preparing the United Nations Demographic Yearbook, for instance, substantial effort was required to ensure that data were properly annotated (via footnotes).

Participants noted that data on population and births would also have to be included to serve as denominators in calculating mortality rates. To make full use of time series of recorded deaths, projected populations would be needed as denominators. The database should make allowance for their inclusion. Meta-information on how such data or other population estimates were obtained would need to be recorded and made available to users. Recalling the activities of UNICEF to make child mortality estimates available, it was suggested that when the Population Division's database went public it should make explicit how United Nations entities collaborated in sharing data.

Participants expressed support for the proposal to collaborate in both developing and populating the database. A pilot database could be developed at the Population Division and shared with collaborators to test the design. It was decided that a working group including the Population and the Statistics Divisions of the United Nations, WHO and the University of California at Berkeley would collaborate in getting the first phases of this project started.

Participants agreed that a web interface would be the best way to share access to the database. Several institutions offered to collaborate in adding data to the database. The World Bank could contribute data derived from its household surveys. WHO would share the data it gathered on mortality statistics.

two major categories with similar patterns of mortality by age and sex. Group 1 reflected mortality in populations with low levels of HIV/AIDS whereas Group 2 included populations with high levels of HIV/AIDS. The average mortality schedules for these two groups were used as standards in a Brass logit relational model to generate model life tables. For each group, derivation of a model implicitly assumed that the populations used to generate the standard were homogeneous.

Mr. Bawah used the examples of Ethiopia and Guinea to show how the models could be used to estimate life tables for national populations. The models provided a means of smoothing estimates derived from observed data and could be used to project mortality by establishing a trend from time-series data

Ms. Meslé focused next on causes of death. Life expectancy in Eastern Europe and the former USSR had largely stagnated since the mid-1960s owing to increasing adult mortality caused mainly by cardiovascular disease and violent deaths. During the 1990s, most Central European countries and the Baltic States had begun to see a reduction of mortality due to cardiovascular disease, but the situation in the Russian Federation and Ukraine was still worsening. It was therefore not easy to foresee what future trends in mortality would be like: a continuation of stagnation or the start of a sustained decline in mortality?

In discussing these findings, participants noted that the causes of the recent improvement in life expectancy in some countries of Central and Eastern Europe were complex. Both improving trends in healthy behaviour and better medical care were contributing to such a trend. Changes in mortality trends in the former German Democratic Republic provided an interesting example of a turnaround driven mostly by improved medical care and access to modern treatments.

Participants agreed that there was considerable uncertainty in projecting mortality for the countries of Central and Eastern Europe because it was not clear whether mortality trends would change in those countries where mortality had largely stagnated or whether the countries whose mortality levels had been declining could sustain that decline over the coming decades. In many of the countries concerned, levels of adult mortality were unusually high relative to levels of child mortality, yielding an atypical age pattern of mortality. Trends in mortality due to cardiovascular disease or in mortality due to violence were expected to determine future changes in overall mortality in many Central and Eastern European countries.

c. Other case studies

Ms. Sawyer presented the cases of three countries China, Mexico and Zimbabwe to illustrate the challenges of producing a best estimate of adult mortality on the basis of multiple but imperfect data sets. She noted that estimates derived from different types of data were not always in agreement and that estimates based on the same data but derived using different methods could differ from one another.

Ms. Sawyer noted that, regarding child mortality, a spline was fitted to all available direct and indirect estimates of child mortality to produce the estimates adopted by the Inter-Agency Group for Mortality Estimation. In fitting the spline, analysts assessed the quality and reliability of the various estimates available and reflected that assessment in terms of weights applied to each estimate during the fitting procedure. With regard to the estimation of adult mortality, where the number of estimates over time was less dense, Ms. Sawyer remarked that fitting procedures to derive trends over time were not being systematically used as yet. To compare the consistency of child mortality estimates with those relative to adult mortality, Ms. Sawyer derived a value of ${}_{45}q_{15}$ from the United Nations one-parameter model life tables using as input the estimated ${}_{5}q_{0}$ as reported by UNICEF and transformed into child mortality by sex by assuming a differential between the sexes.

gramme (DHS) had collected information on sibling survival and censuses and surveys had recorded data on parental survival. The estimates derived from those sources indicated without doubt that adult mortality had been increasing since the late 1980s. However, estimates of ${}_{45}q_{15}$ derived from different sources for the same period differed by wide margins.

During the discussion, participants remarked that the procedure used to transform child mortality estimates into an indicator of adult mortality was useful in exploring the quality and consistency of the estimates at hand. In that respect, lack of a consistent and validated time series of child mortality estimates by sex for each country was a drawback. Noting that sex differentials in childhood mortality varied across countries and over time, participants underscored the need to derive consistent time series of child mortality estimates by sex.

I. Approaches to modelling AIDS mortality

Mr. Vallin expressed support for the proposal of compiling all available information and promoting the sharing of such information. He encouraged the development of new methods of mortality estimation and the further refinement of older methods. Demographic models had to be tested against reality. He suggested that sub-national data could be used to that end. Mr. Vallin commended the work of the different entities producing estimates of mortality and underscored the usefulness of sharing results and cooperating in comparing data and methods to ascertain the causes of any discrepancies that might arise. If absolute consistency among the sets of estimates produced by different entities could not be reached, a useful approach might be to find ways of communicating systematically the degree of uncertainty to which different estimates were subject.

To conclude, Mr. Wilmoth summarized the expected follow-up to the proposals made at the meeting by focusing on three main issues:

Database development. Mr. Wilmoth underscored the importance of this activity to set the basis for the systematic derivation and assessment of mortality estimates. He recalled that a sub-group constituted by the United Nations Population Division, the United Nations Statistics Division, WHO and the University of California at Berkeley would collaborate to develop a pilot version of the database, test it and develop a protocol for data exchange. This pilot version would be presented and discussed at the next coordination meeting on mortality.

Further development and testing of estimation methods. Mr. Wilmoth emphasized the need for further work on evaluating or developing indirect methods to estimate adult mortality, noting that there was mixed evidence about the reliability of some methods when their underlying assumptions were violated. The next coordination meeting on adult mortality might address those issues morty

m1263 evidence of ( )]-5tingfor (ation)-2(r)(at(rtalit))-8( )]TJ0.000
t0.361ess thatpartn6(b 7.98 0 0 12 -1.1598 2147Td50nd disc(hipsg

ORGANIZATION OF WORK *(CONTINUED)*

Friday, 27 October 2006

Morning session: *8:30-12:00*

6. ESTIMATING ADULT MORTALITY USING INCOMPLETE VITAL REGISTRATION DATA

a) Evaluating the validity of indirect estimates of adult mortality: Mr. Ken Hill
b) Simulation analysis of death distribution methods (DDM): Ms. Danzhen You
c) Comparison of data sources for DDM estimation: Ms. Mila Andreeva
d) Toward a methods protocol for DDM estimation: Mr. John Wilmoth

Morning break: *10:30-10:45*

7. DATABASE DEVELOPMENT

a) Database design and data processing:

UN/POP/MORT-EGM/2006/INF.2

1 November 2006

English only

UNITED NATIONS EXPERT GROUP MEETING ON
CURRENT ISSUES IN THE ESTIMATION OF ADULT MORTALITY
United Nations Secretariat
Department of Economic and Social Affairs
Population Division
New York, 26-27 October 2006

LIST OF PARTICIPANTS

Mrs. Lioudmila ANDREEVA
Research Specialist
Rockefeller University
Laboratory of Populations
1230 York Avenue
New York, NY 10021
Tel: 1-212-327-7536
Fax: 1-212-327-7974
Email: mandreeva@rockefeller.edu

Mr. Ayaga A. BAWAH
Senior Research Associate
INDEPTH Network
11 Mensah Wood Street, East Legon
P. O. Box KD 213, Kanda
Accra, Ghana.
Tel/fax: 233-21-519394
Email: bawah.ayaga@indepth-network.org

Mr. Griffith FEENEY
Consultant
UN Population Division
9 Fairview Road
Scarsdale, NY 10583
Tel: 914-715-3224
Email: griff@griffithfeeney.com

Ms. Dana A. GLEI
Research Demographer
University of California, Berkeley
5985 San Aleso Ct.
Santa Rosa, CA 95409-3912
Tel/Fax: 707-539-5592
Email: danaglei@demog.berkeley.edu

Mr. Ken HILL
Professor
Harvard Center for Population
and Development Studies
9 Bow Street
Cambridge, MA 02138
Tel: 617-496-6708
Fax: 617-495-5418
Email: kenneth_hill@harvard.edu

Ms. France MESLÉ
Director of Research
Institut national d'études démographiques
133 bd. Davout
75890 Paris cedex 20
Tel: 33 1 56 06 21 43
Fax: 33 1 56 06 21 94
Email: mesle@ined.fr

Mr. Vladimir CANUDAS-ROMO
Researcher
Department of Demography
University of California, Berkeley
2232 Piedmont Ave.
University of California, Berkeley
Berkeley, CA 94720-2120
Tel: 510-642-9471
Fax: 510-643-8558
Email: canudas@demog.berkeley.edu

Mr. Jacques VALLIN
Director of Research
Institut national d'études démographiques
133 bd. Davout
75980 Paris cedex 20
Tel: 33 1 56 06 21 06
Fax: 33 1 56 06 21 94
Email: vaoil,U10.98 0 0 10.98 300.481ed.fMCID 5 BDC BT/TT0 1 Tf20 Tc 10.98 0 0 10.98 72 5872 595.8013 Tm[(Em)9

United Nations Children's Fund

Mr. Neff WALKER
Senior Programme Officer
Strategic Information Section
Division of Policy and Planning
3 UN Plaza, H-0488
New York, NY 10017
Tel: 212-326-7258
Fax: 212-735-4411
Email: walkern@unicef.org

Pan American Health Organization/WHO

Mr. John SILVI
Statistician
Unit of Health Analysis and Statistics
525 Twenty-third Street N.W.
Washington, DC. 20037
Tel: 202-974-3141
Fax: 202-974-3674
Email: silvijhn@paho.org

DEPARTMENT OF ECONOMIC AND SOCIAL AFFAIRS

Statistics Division

Mr. Srdjan MRKIC
Acting Chief
Demographic Statistics Section
Two UN Plaza, DC2-1556
New York, NY, 10017
Tel: 212-963-4940
Fax: 212-963-1940
Email: mrkic@un.org

Mrs. Haoyi CHEN
Statistician
Demographic Statistics Section
Two UN Plaza, DC2-1536
New York, NY, 10017
Tel: 212-963-4375
Fax: 212-963-1940
Email: chen9@un.org

Ms. Sabine WARSCHBURGER
Statistician
Demographic Statistics Section
Two UN Plaza, DC2-1562
New York, NY, 10017
Tel: 212-963-4297
Fax: 212-963-1940
Email: warschburger@un.org

Mr. Gerhard HEILIG
Chief
Population Estimates and Projections Section
Two UN Plaza, DC2-1920
New York, NY, 10017
Tel: 212-963-4531
Fax: 212-963-2147
Email: heilig@un.org

Ms. Cheryl SAWYER
Population Affairs Officer
Mortality Section
Two UN Plaza, DC2-1986
New York, NY, 10017
Tel: 212-963-3219
Fax: 212-963-2147
Email: sawyerc@un.org

Mr. François PELLETIER
Population Affairs Officer
Population Estimates and Projection Section
Two UN Plaza, DC2-1914
New York, NY, 10017
Tel: 212-963-3213
Fax: 212-963-2147
Email: pelletierf@un.org

Mr. Kirill ANDREEV
Population Affairs Officer
Mortality Section
Two UN Plaza, DC2-1984
New York, NY, 10017
Tel: 212-917-9352
Fax: 212-963-2147
Email: andreev@un.org